# First Pass Success Rates of Out-Of-Hospital Advanced Airway Management in Adults and Children



Jeffrey L. Jarvis, MD, MS, EMT-P<sup>1</sup>, David Wampler, PhD, EMT-P<sup>2</sup>, Henry Wang, MD, MS<sup>3</sup>

<sup>1</sup>Williamson County EMS, <sup>2</sup>UT Health Science Center – San Antonio, <sup>3</sup>UT Health Science Center - Houston



#### Introduction

Prehospital advanced airway management (AAM, including endotracheal intubation (ETI) and supraglottic airway insertion (SGA)) of children is difficult. Multiple AAM attempts are associated with increased adverse events. We sought to compare AAM first-pass success (FPS) rates between adults and children in a national cohort of EMS agencies.

# **Methods**

We used 2017 clinical data from ESO, a national electronic health record system encompassing over 2,000 EMS agencies. We included all patients receiving any AAM attempts. FPS success was self-reported. Using multivariable logistic regression, we compared the odds of ETI FPS between adults (age >=14 years) and children (age <14 years), adjusting for gender, ethnicity, primary impression, and drug facilitation. We repeated the analysis for SGA FPS. Statistics were performed using R. Approval was waived by the Baylor Scott & White IRB.

#### Results

During the one year period, 731 EMS agencies attempted AAM on 29,369 patients (median 18 per agency [IQR 6, 43]).


## Conclusions

ETI FPS is lower in children than adults.

SGA FPS does not differ between children and adults. It may be reasonable to adopt an SGA first approach in pediatric patients.

Characteristics of Data Set by Adult versus Pediatric								
		Adult	Pedi					
n		28846	523					
		65.06	2.22					
Age (median [IQR])		[51.66, 76.38]	[1.19, 6.74]					
Identity (%)	Asian	231 (0.8)	6 (1.1)					
	Black	5091 (17.6)	119 (22.8)					
	Hispanic	610 (2.1)	23 (4.4)					
	Native Am	80 (0.3)	5 (1.0)					
	Not Known	4090 (14.2)	96 (18.4)					
	White	18744 (65.0)	274 (52.4)					
Gender (%)	Female	11209 (38.9)	217 (41.5)					
	Male	17464 (60.5)	305 (58.3)					
	Not Reported	173 (0.6)	1 (0.2)					
		40406 (67.2)	254/677\					
Primary Imp. (%)	Cardiac Arrest	19406 (67.3)	354 (67.7)					
	Medical	7297 (25.3)	100 (19.1)					
	Trauma	2143 (7.4)	69 (13.2)					
Airway Method (%)	ETI	22049 (76.4)	471 (90.1)					
	SGA	6797 (23.6)	52 (9.9)					
Drug Assisted (a)	Not a DAI	22126 (90.2)	4EC (07.2)					
Drug Assisted (%)	Not a DAI	23136 (80.2)	456 (87.2)					
	RSI SAI	4063 (14.1)	50 (9.6)					
		1647 (5.7)	17 (3.3)					
	911 Response	28,562 (99.0)	513 (98.3)					
Dun Tun o (o)	Interfacility	201 (0.7)	0 /1 7)					
Run Type (%)	Transfer	201 (0.7)	9 (1.7)					
	Mutual Aid	74 (0.3)	0 (0.0)					
	Other	9 (0.0)	0 (0.0)					

Adult and Pediatric FPS with ETI and SGA								
		ETI FPS	Adjusted OR		SGA FPS	Adjusted OR		
	ETI (n)	Rate (%)	(95%CI)	SGA (n)	Rate (%)	(95%CI)		
Adult								
(referent)	22,049	72.7%		6,797	89.8%			
			0.6			0.6		
			(0.5 to 0.7)			(0.3 to 1.4)		
Pedi	471	58.5%	p < 0.001	52	84.6%	p = 0.24		
Age								
Groups								
			0.44			0.2		
<1	78	52.6%	(0.2 to 0.7)	6	66.7%	(0.04 to 1.7)		
			0.5					
1 to <6	274	55.8%	(0.4 to 0.6)	20	100%	-		
			0.7			0.2		
6 to <10	52	63.5%	(0.4 to 1.3)	7	57.1%	(0.04 to 0.9)		
			0.9			0.4		
10 to <14	103	69.9%	(0.6 to 1.4)	23	78.3%	(0.2 to 1.2)		
=>14								
(referent)	22,013	72.7%	p < 0.001	6,793	89.9%	p = 0.44		



## Acknowledgements

The authors wish to express their appreciate to ESO for its assistance with the data.